SYMAN: a SYMmetry ANalyzer

Ari D. Gross and Terrance E. Boult
Department of Computer Science, Columbia University NYC NY 10027.
ari@cs.columbia.edu, boult@cs.columbia.edu

Abstract

Symmetry is pervasive in both man-made objects and nature. Since symmetries project to skew symmetries, finding axes of skew symmetry is an important vision task. This paper describes the construction of a symmetry analyzer. Examples using SYMAN on both real and synthetic images are shown. We motivate SYMAN's combination of both global and local methods. We show how to derive a global analytic solution for the skew axes when the degree of skew symmetry is known. We also present a new local tangent-based algorithm which has advantages over previous methods.

1 Introduction

A reflective symmetric planar figure projects, under orthography and weak perspective, to a skew symmetric contour. *Skew symmetry* is defined in [4] as a symmetry at a fixed angle to the axis. An intensity image, with detected edges overlayed, is shown in figure 1 (left). The right side of figure 1 displays SYMAN's recovered axes from the noisy hand-segmented contour shown.

Previous work on finding axes of skew symmetry can be divided into two classes: local and global. Local techniques compare pairs of points. Such points, if they are pairwise skew symmetric, satisfy necessary constraints [3], [6], [7]. Local methods can be robust under occlusion but, because of the number of pairs, are computationally expensive. For unoccluded contours, global moment-based techniques can be more efficient. A moment-based algorithm for finding axes of skew symmetry was proposed in [2] but requires searching two 1D spaces using a heuristic symmetry evaluator. A hough-based method is given in [8] but no results are presented.

The next section presents some examples of the SYMAN's performance. In section 3 we show how to derive an analytic solution for the axes of skewed symmetry, given the number of solutions. In section 4, we present a local tangent-based technique and discuss its advantages. Section 5 discusses the tradeoffs between local and global techniques for recovering axes of skew symmetry. The paper ends with some conclusions and a very brief discussion of future work.



Figure 1: PUMA intensity image with overlayed edges (left) and axes of skew symmetry recovered by SYMAN for the hand-segmented contour shown (right).

2 SYMAN Examples

We now present a few examples of SYMAN's capabilities, some of which are discussed elsewhere in the paper. For more details on SYMAN's capabilities see [3]. Figure 2 is a die image with axes of skew symmetry recovered by SYMAN using both local and global methods. A small camera-imaged contour of a figure from [5] is shown in figure 3, on its right is the deskewed, derotated form recovered by SYMAN. A leaf image, with overlayed edges, is shown in figure 4. The right side of that figure shows the derotated, deskewed contour recovered by SYMAN.

3 Deriving an Analytic Solution for Axes of Skew Symmetry

Given a contour C assume its degree of skew symmetry, $deg(C)$ = number of valid skew symmetry axes, can be determined *á priori*, see [3]. Knowing $deg(C)$ we want to recover all of the skew axes. Assume, for now, that $deg(C) = 3$. We start by computing the four 3^{rd} order moments of C. As noted in [2], any weak perspective or orthographic projection of a symmetric planar figure S can be represented as a image-plane skew followed by a rotation. Without loss of generality, let S be symmetric about the y-axis. The 3^{rd} order moments of C are functions of the 3^{rd} order moments of S, a rotation parameter α and a skew parameter β. Since S is symmetric, the $x^{odd}y^j$ moments are zero. Thus, the four 3^{rd} order moments for contour C are functions of α, β, and two 3^{rd} order moments of S. Using elimination theory on the four equations derived for the 3^{rd} order moments



Figure 2: Die image due to Glenn Healey via Jean Ponce. Results using local tangent-based method (top) and moment-based method (botom). Runtimes per contour were 30-60 sec. and 1 sec., respectively.



Figure 3: On the left is a skewed and rotated edge image (obtained pointing a camera at a figure in [5]) and on the right is a solution recovered by SYMAN.



Figure 4: Intensity image of a leaf, with hand-segmented edges of the leaf overlayed (left) and the derotated, deskewed leaf contour recovered by SYMAN.

of C, we can derive a resultant cubic univariate polynomial in rotation parameter α. For each α solution, a corresponding β solution can be obtained.

For any degree of skew symmetry $n = deg(C)$, an analogous procedure is used that relies on the n^{th} order moments of C and S. The n^{th} order polynomial can be precomputed using elimination theory [1]. If we know $deg(C) = n$, the n^{th} order contour moments of C are computed, and the associated n^{th}order polynomial equation in α is solved. SYMAN has the precomputed form of these polynomials through degree 10.

Figure 3 shows a skew symmetric contour c_M, where the original symmetric figure is from [5]. SYMAN hypothesized that the contour c_M is 2-fold symmetric (see [3]), that $deg(c_M) = 4$, and then solved for the 4 sets of axes using the above method. One of the 4 recovered solutions, displayed as a deskewed and derotated contour, is shown in figures 3. When presented with the contour of a skewed and rotated 9-gon and given $deg(c_9) = 9$, SYMAN used the moment-based method to approximately recover all 9 solutions (see [3]).*

4 Our Local Tangent-Based Method

Several local methods for finding axes of skew symmetry have been proposed in the literature. These are essentially algorithms constructed around a constraint between skew symmetric pairs of contour points. We propose an algorithm based on a new skew symmetry constraint and contrast it with two existing local methods [6],[7]. SYMAN has running versions of all three methods.

Theorem 1 *Skew Symmetry Tangent Theorem: If a pair of points P_+ and P_- are skew symmetric, then they satisfy a pairwise constraint given by $2 \cdot \cot\beta = \cot\theta_- + \cot\theta_+$, where β is the angle of skew and θ_+ (θ_-) the angle between the transverse axis and the tangent at P_+ (respectively P_-).*

The local tangent-based algorithm considers every pair of contour points, letting each pair vote for a set of

*While the method uses high order moments, something about the ratio of moments that it uses appears to make it reasonably robust. A formal investigation is underway.